Efficacy and safety of erbium-doped yttrium aluminium garnet fractional resurfacing laser for treatment of facial acne scars

Geethu F Alapatt^{1*}, Ramesh Bhat M², Sukumar D³, Meryl Sonia Rebello⁴

^{1,4}PG Student, ²Professor and HOD, ³Professor, Department of Dermatology, Father Muller Medical College, Mangalore-575002, Karnataka, INDIA.

Email: geethu.f.alapatt@gmail.com

Abstract Background: Acne scarring has lifelong sequelae and are extremely disturbing to patients, both physically and psychologically. The use of ablative fractional lasers for the treatment of acne scars is becoming increasingly popular. But studies regarding the efficacy and safety of the same in the Indian skin is limited. Objective: To evaluate the efficacy and safety of Erbium-doped Yttrium Aluminium Garnet (Er: YAG) 2940 nm fractional laser resurfacing in the treatment of acne scars in 10 patients at a tertiary care teaching hospital Methodology: All 10 patients received four treatment sessions with Er: YAG fractional laser at 1-month interval. Subjective assessment in percentage of improvement was documented after four sessions. Subjects were instructed to report any cutaneous side-effects including erythema, oozing, crusting, dyschromia, scarring or secondary infection and about interference with daily activities in the post-treatment period. Photographs were taken before each treatment session and 1 month after the final session. A clinical assessment by comparing the photographs was done. Patient's satisfaction of improvement was noted at the end of four sessions. Conclusion: Er- YAG laser resurfacing is both effective and safe in the treatment for acne scars. Keywords: Ablative, acne scars, efficacy, fractional, laser, safety.

*Address for Correspondence:

Dr. Geethu F. Alapatt, PG Student, Department of Dermatology, Father Muller Medical College, Mangalore-575002, Karnataka, INDIA. **Email:** <u>geethu.f.alapatt@gmail.com</u>

Received Date: 10/08/2015 Revised Date: 01/09/2015 Accepted Date: 28/10/2015

Access this article online		
Quick Response Code:	Website: <u>www.statperson.com</u>	
	DOI: 01 November 2015	

INTRODUCTION

Acne scars are seen after inflammatory acne and is caused by the destruction of collagen.¹ They have a lifelong sequelae and are extremely disturbing to patients, both physically and psychologically.² They are very difficult to treat.³ A number of treatment modalities are available which includes dermabrasion, subcision, chemical peeling, fillers and punch technique but these are not very effective. Lasers have been increasingly used for the treatment of acne scars. Ablative lasers show good clinical improvement but have a lot of adverse effects like dyspigmentation and post- procedure erythema, especially in people with dark skin. Non ablative lasers do not cause many adverse effects but do not show much clinical improvement. To overcome the drawbacks of the ablative and non ablative lasers, fractional laser resurfacing techniques were developed.⁴ However studies on the efficacy and safety profiles of this type of laser in Indian skin is lacking.

MATERIALS AND METHODS

This was a prospective study performed in a tertiary hospital from March 2014 to April 2015 and included patients with clinically diagnosed scarring secondary to acne. Ethical clearance was obtained from the ethical committee of the institution. Patients who were 18 yrs of age and above with presence of acne scars were included. Patients with known photosensitivity, pregnancy, lactation, keloidal tendency, use of oral retinoids in last 6 months, infections like herpes labialis, active acute illness, and unrealistic expectations were excluded. A written informed consent was taken from each patient enrolled in this study. A detailed history and clinical

How to site this article: Geethu F Alapatt, Ramesh Bhat M, Sukumar D, Meryl Sonia Rebello. Efficacy and safety of erbium-doped yttrium aluminium garnet fractional resurfacing laser for treatment of facial acne scars. *International Journal of Recent Trends in Science and Technology* November 2015; 17(1): 01-04 <u>http://www.statperson.com</u> (accessed 02 November 2015).

examination was carried out. Area to be treated was cleansed with a mild cleanser and normal saline. Topical anesthesia with 2.5% lidocaine and 2.5% prilocaine cream (EMLA) was applied 60 minutes before treatment. They were treated with fractional ablative resurfacing module using 2940 nm Er: YAG handpiece. Parameters for each sitting were kept constant in all patients. Long pulse mode was used for all four sessions. 3 passes in vertical, horizontal, and oblique directions with up to 50% overlapping was given. Ice packs were applied for at least 10 minutes after procedure. Strict photoprotection and topical steroid cream was recommended 3 days postprocedure. Photographic documentation using identical camera settings and patient positioning was obtained at baseline and before each session. Patient satisfaction was graded after four treatment sessions as

- Grade 1: <25%, minimal to no improvement
- Grade 2: 26-50%, moderate improvement;
- Grade 3: 51-75% marked improvement

Grade 4: >75% almost complete clearance of scarring

Subjects were asked to report any side effects like pain, erythema, oozing, edema, crusting, scarring, hyperpigmentation or secondary infection during and after the procedure in the treatment period of four months. They were asked about interference with daily activities in the post treatment period. Data was analyzed using percentage and frequency,

RESULTS

A total of 10 patients with acne scars participated in the study. The skin phototype of all these patients were IV-V.



Figure 1: Before treatment

The study comprised of 6 males and 4 females and all were in the age group of 18-35yrs. Most of the patients (50%) reported a marked improvement in the acne scars at the end of 4 sittings. Moderate improvement was seen in 40% of the patients. Only 10% that is one out of the 10 patients reported no improvement at the end of 4 sittings. The most common adverse effects were pain during the procedure and erythema following the procedure. Most of the patients (80%) reported erythema following the procedure which lasted for 1-2 days. Only 1 patient had prolonged erythema which lasted 4 days. One out of the ten patiens did not have any ervthema following the procedure. Some amount of pain was complained by 80% of the patients during the procedure. One out of them had severe pain. . No other side effects like oozing, crusting, dyschromia, scarring, secondary infection or exacerbation of acne lesions were reported.

Table 1: Grading of improvement by patient and side effects

<u> </u>	/ 1
Grade of	Side effects
improvement	Side effects
3	Erythema,
3	Erythema, severe pain
2	Eythema, mild pain
3	Erythema, mild pain
2	Mild pain
2	Erythema Mild pain
2	Erythema Mild pain
3	Erythema Severe pain
2	Prolonged erythema Mild
5	pain
1	Erythema
	Grade of improvement 3 2 3 2 2 2 2 3 3 3 3 1



Figure 2: After four sessions



Figure 3: Before treatment



Figure 4: After four sessions



Figure 5: Before treatment



Figure 6: After four sessions



Figure 7: Before treatment

DISCUSSION

This study demonstrated the efficacy and safety of Erbium-doped Yttrium Aluminium Garnet fractional resurfacing laser for treatment of facial acne scars. Acne scars are the result of impaired resolution following inflammatory acne. They are classified as ice-pick scars, rolling scars and box scars on the basis of the shape and depth of these scars.⁴ The use of ablative carbon dioxide lasers dates back to 1980s.⁵ These are based on the theory of selective photothermolysis.⁶ These lasers though had a good efficacy did not have a good safety profile. Thus non-ablative lasers were developed with better safety profile but their efficacy was unpredictable.⁷ Manstein and Colleagues devised Fractional photothermolysis (FP) in 2004.⁸ But the results of these lasers were not satisfactory. Thus lasers with the combination of FP



Figure 8: After treatment

theory and ablative lasers technology were devised. FP targets water as a chromophore. Epidermal keratinocytes, collagen and blood vessels are the water rich targets in skin. Ablative laser resurfacing using CO2 or Er: YAG lasers has shown an efficacy of 25% to 90% for the treatment of acne scars and is considered the gold standard. But post procedure complications like erythema, infection, scarring and pigmentary changes especially in dark skin like the Indian skin were common. ⁴ A study done by Nirmal *et al* has reported ablative fractional photothermolysis as an effective and safe treatment for atrophic acne scars in Indian skin.⁷ A study done by Ahmad et al on the use of CO2 fractional laser in the treatment of acne scars showed 25% to75% improvement in most patients. Another study done by Saryazdi et al showed an objective and subjective improvement of about 20-70% and 30-70%, respectively, without any erythema, permanent hyperpigmentation and other adverse effects.9 Our study also showed similar findings. This study demonstrated that at the end of four sitting 50% of the patients reported a marked improvement in the acne scars. Moderate improvement was seen in 40% of the patients. Only 10% that is one out of the 10 patients reported no improvement in acne scarring at the end of 4 sittings. All the patients reported improvement in skin texture. The most common adverse effects were pain during the procedure and erythema following the procedure. Most of the patients (80%) reported erythema following the procedure which lasted for 1-2 days. Only 1 patient had prolonged erythema which lasted 4 days. One out of the ten patiens did not have any erythema following the procedure. Some amount of pain was complained by 80% of the patients during the procedure. One out of them had severe pain. Two out of the ten patient did not experience any pain during the procedure. No other side effects like oozing, crusting, dyschromia, scarring, secondary infection or exacerbation of acne lesions were reported.

CONCLUSION

The results of this study suggest that ER: YAG fractional laser may be a useful tool in treating acne scarring with minimal side effects.

REFERENCES

1. Holland DB, Jeremy AH, Roberts SG, Seukeran DC, Layton AM, Cunliffe WJ. Inflammation in acne scarring:

A comparison of the responses in lesions from patients prone and not prone to scar.Br J Dermatol 2004; 150:72-81.

- Lee SJ, Kang JM, Chung WS, Kim YK, Kim HS. Ablative non-fractional lasers for atrophic facial acne scars : a new modality of erbium: YAG laser resurfacing in Asians. Lasers Med Sci 2014;29:615-9
- Holland DB, Jeremy AH, Roberts SG, Seukeran DC, Layton AM, Cunliffe WJ. Inflammation in acne scarring: A comparison of the responses in lesions from patients prone and not prone to scar.Br J Dermatol 2004; 150:72-81.
- Ahmad TJ, Muzaffar F, Nabi H, Malik S, Noreen A, Rabiya Hayat. Efficacy and safety of ablative fractional carbon dioxide laser for acne scars Journal of Pakistan Association of Dermatologists 2012;22:41-4
- 5. Brightman LA, Brauer JA, Anolik R, Weiss E, Karen J, Chapas A, *et al.* Ablative and fractional ablative lasers. Dermatol Clin 2009; 27:479-89.
- Anderson RR, Parrish JA. Selective photothermolysis: Precise microsurgery by selective absorption of pulsed radiation. Science 1983; 220:524-7.
- Nirmal B, Pai SB, Sripathi H, Rao R, Prabhu S, Kudur MH, *et al.* Efficacy and safety of Erbium-doped Yttrium Aluminium Garnet fractional resurfacing laser for treatment of facial acne scars. Indian J Dermatol Venereol Leprol 2013; 79:193-8.
- Manstein D, Herron GS, Sink RK, Tanner H, Anderson RR. Fractional photothermolysis: A new concept for cutaneous remodeling using microscopic patterns of thermal injury. Lasers Surg Med 2004; 34:426-38.
- Saryazdi S, Mohebbi A. Evaluation of Efficacy of Fractional CO2 Laser in Acne Scar: J Lasers Med Sci 2012; 3:56-60

Source of Support: None Declared Conflict of Interest: None Declared