

A clinico-demographic study of reproductive tract infections at tertiary health care center

Birajdar Annasaheb Gulabrao^{1*}, More N V², Suwarna Dahitankar³

¹Assistant Professor, Associate Professor, Department of OBGY, Government Medical College , Latur, Maharashtra, INDIA.

²Ex Assistant Professor, SRTR Government Medical College, Ambajogai, Beed, Maharashtra, INDIA.

Email: drannasaheb83@gmail.com

Abstract

Introduction: the maximum cases of Bacterial vaginosis, Trichomonasvaginalis and Candidiasis were in the age group of 20-29 yrs. and least in the age group of ≤ 19 yrs. As parity increases, the no. of cases of each one become less and most of the cases were having parity 1 to 2. **Aims and Objectives:** To Study Clinico -Demographic profile of patients with Reproductive Tract Infections at tertiary Health care center. **Methodology:** Patients attending tertiary health care center with complaint of abnormal vaginal discharge of age group 15 to 40 years were selected for the study. This study was conducted at tertiary health care centre from January 2009 to September 2010. During the study period the total number of gynaec O.P.D attendees were 19,270. Presuming the equal distribution of cases on all days the no. of attendees (symptomatic and asymptomatic) with findings of RTI were 6,423. Data presented in tables and in percentages. **Result:** Vaginal discharge was the commonest complaint among the study group. Prevalence increased with increase in Parity. RTI was more among the Muslims. Prevalence was higher among non-working women. Prevalence of RTI cases was higher in Illiterate. The prevalence of RTI was highest in non-users of Contraceptives (41.26%) as compared to other groups; it was lowest in the women who were dependent on their spouse for the use of condom as a contraceptive. Out of 64 cases of Bacterial vaginosis, 43.75% complained of vaginal discharge, 37.5% complained of pain in abdomen, 15.62% patients complained of backache, 20.31% complained of itching at genital area, 23.43% complained of increased frequency of micturition and 10.93% complained of dyspareunia. Among 40 cases of Trichomonasvaginalis, vaginal discharge was present in 40% cases, pain in abdomen in 20%, backache in 22.5%, Itching in 45%, increased frequency of micturition in 27.5%, dyspareunia in 12.5% of cases. Among 36 cases of Candidiasis vaginal discharge was present in 59.45% cases, backache in 37.83%, pain in abdomen in 40.54%, itching at genital area in 51.35%, increased frequency of micturition in 29.72 % and dyspareunia in 24.32%. **Conclusion:** It can be concluded from study that Majority of cases with suggestive symptoms of RTI were in the age group of 20-29 years, asymptomatic cases were highest in the parity group > 4 . Prevalence of RTI cases was high among Muslims nonworking (housewife), in illiterate and in non-users of contraceptives. In the study population cases were having multiple symptoms of which in overall vaginal discharge was the most common complaint.

Keywords: Reproductive Tract Infections, Trichomonasvaginalis, Bacterial vaginosis, Candidiasis.

*Address for Correspondence:

Dr. Birajdar Annasaheb Gulabrao, Assistant Professor, Associate Professor, Department of OBGY, Government Medical College , Latur, Maharashtra, INDIA.

Email: drannasaheb83@gmail.com

Received Date: 10/09/2016 Revised Date: 15/10/2016 Accepted Date: 12/11/2016

Access this article online

Quick Response Code:



Website:
www.statperson.com

DOI: 15 November
2016

INTRODUCTION

Vaginal discharge is one of the main complaints of the patients attending gynaecology O.P.D. The term 'Leucorrhoea' should be used to denote the normal vaginal secretions that are increased in amount. An increase in the normal vaginal secretions develops physiologically at puberty, during pregnancy and at ovulation. An abnormal vaginal discharge causes much more morbidity in the form of physical discomfort and psychological distress. So it is necessary to diagnose & treat it. The normal vaginal flora is mostly aerobic but also contains anaerobic bacteria. Several factors influence

the vaginal flora including Age, P^H, Sexual activity, Phase of menstrual cycle, Use of antibiotics & Any Invasive Procedure. The most frequent aerobic bacteria are Lactobacillus, Gardnerellavaginalis, Staphylococcus, E. coli. Anaerobes include Peptostreptococci, Peptococci, Prevotella etc. P^H of the normal vagina is 4.5, which is maintained by the production of lactic acid. An alteration in the normal vaginal flora is an important factor predisposing for vaginal infection. Any vaginal discharge that is frankly purulent & contains pus cells should be considered due to specific vaginal infection. Symptoms such as vaginal discharge, itching, irritation are common reasons for consulting the general practitioner. Sexually transmitted infections (STIs) are the infections caused by germs such as bacteria, viruses or protozoa that are passed from one person to another mainly through sexual contact. STIs are a part of a broader group of infections known as reproductive tract infections (RTIs) that includes all the infections of the reproductive tract, including those not caused by sexual contact. The term "STD" denotes the full-blown disease but does not represent "asymptomatic infections". STI is a better description as it includes all infections, both symptomatic and asymptomatic. Reproductive tract infections (RTIs), including both sexually transmitted infections (STIs) and non-sexually transmitted infections (non-STIs) of the reproductive tract are responsible for major ill-health throughout the world.¹ World Health Organization estimates that each year there are over 340 million new cases of sexually transmitted infections in which 75–85% occur in developing countries. In India alone, 40 million new cases emerge each year.² A majorities of women continue to suffer from RTIs leading to complications like pelvic inflammatory disease (PID), infertility, cervical cancer, postabortal, and puerperal sepsis, chronic pelvic pain, and ectopic pregnancy. RTIs in many cases are asymptomatic among women, making their detection and diagnosis difficult.³ Reproductive tract infections (RTIs) including sexually transmitted infections (STIs) are recognized as public health problem and rank second as the cause of healthy life lost among women of reproductive age after maternal morbidity and mortality in developing countries. Emerging epidemics of Acquired immunodeficiency syndrome (AIDS) and identification of STIs as a co-factor in its acquisition have made the control of these infections, one of the strategies imperative to decrease the transmission of HIV/AIDS. World over 340 million people are affected by STIs, out of which 30 million reside in India⁴.

MATERIALS AND METHODS

Patients attending tertiary health care center with complaint of abnormal vaginal discharge of age group 15

to 40 years were selected for the study. While Patients who are menstruating, on any antibiotics or vaginal medication during period of 7 days, Patients who are having genital malignancy or fibroid uterus were excluded from the study. This study was conducted at tertiary health care centre from January 2009 to September 2010. During the study period the total number of gynaec O.P.D attendees was 19,270. However the number of cases in the study group was examined by only one person and hence the percentage related to symptomatology cannot be treated as accurate. Presuming the equal distribution of cases on all days the no. of attendees (symptomatic & asymptomatic) with findings of RTI were 6,423. A detail history with special reference to dysuria, vaginal discharge, pelvic pain and menstrual irregularities were noted. General, physical and local examination of the genitalia was carried out and clinical diagnosis was kept. Necessary tests, vaginal swabs were sent for routine, microscopic examination and also for the culture & sensitivity testing to the department of microbiology.

RESULT

Table 1: Age wise distribution of cases

Age group (Yrs)	Bacterial Vaginosis 64	Trichomonas Vaginalis 40	Candidiasis 37
≤ 19	5 (8)	1 (1)	2 (5)
20-29	46 (72)	28 (71)	21 (57)
≥ 30	13 (20)	11 (28)	14 (38)

(Figures in line parenthesis show percentages.)

The present study was done in the age group of 15 to 40 yrs. Above table shows the age wise distribution of clinically diagnosed cases. Total no. of cases of Bacterial vaginosis in the age group ≤ 19 yrs were 5 (8%), in the age group between 20-29 were 46 (72%) and above 30 yrs were 13 (20%). Of Trichomonasvaginalis, in the age group ≤ 19 yrs total cases were 1 (1%), in the age group between 20-29 yrs were 28 (71%) and above 30 yrs were 11 (28%), while that of Candidiasis in the age group ≤ 19 yrs total cases were 2 (5%), in the age group between 20-29 yrs were 21 (57%) and above 30 yrs were 14 (38%). It was observed that the maximum cases of Bacterial vaginosis, Trichomonasvaginalis and Candidiasis were in the age group of 20-29 yrs and least in the age group of ≤ 19 yrs.

Table 2: Parity wise distribution of cases

Parity	Bacterial Vaginosis 64	Trichomonas Vaginalis 40	Candidiasis 37
Nulliparous	16(25)	14(35)	12(32.43)
1-2	24(37.5)	20(50)	16(43.24)
3-4	15(23.43)	5(12.5)	5(13.5)
> 4	9(14)	1(2.5)	4(10.81)

(Figures in line parenthesis show percentages)

Above table shows parity wise distribution of cases of Bacterial vaginosis. Out of 64 cases of Bacterial Vaginosis nulliparous cases were 16 (25%), 24 (37.5%) cases were having parity of 1-2, 15 (23.43%) cases were having parity of 3-4 and 9 (14%) cases were having parity of >4. In case of Trichomonas Vaginalis, the no. of nulliparous cases were 14 (35%), 20 (50%) cases were having with parity of 1-2, 5 (12.5%) cases were having parity of 3-4 and 1 (2.5%) cases were having parity of >4. Out of 37 cases of Candidiasis the no. of nulliparous cases were 12 (32.43%), 16 (43.24%) cases were with parity of 1-2, 5 (13.5%) cases were having parity of 3-4 and 4 (10.81%) cases were with parity of >4. This shows that as parity increases the no. of cases of each one become less and most of the cases were having parity 1 to 2.

Table 3: Symptom wise distribution of cases

Symptom	No. of Cases	Percentages (%)
Vaginal Discharge	89	44.5
Pain in Abdomen	67	33.5
Backache	54	27
Itching	76	38
Increased Frequency of Micturition	53	26.5
Dyspareunia	30	15

Above table shows that out of 200 cases examined 89 (44.5%) patients complained of foul smelling vaginal discharge, 67 (33.5%) complained of pain in abdomen, 54 (27%) of backache, 76 (38%) of itching, 53 (26.5%) of Increased frequency of micturition and 30 (15%) complained of dyspareunia. Therefore it appears that vaginal discharge was the commonest complaint among the study group.

Table 4: Distribution of cases according to religion and suggestive findings of RTI

Religion	Total No. of cases 200	No. of patients	
		Symptomatic 141(70.5)	Asymptomatic 59(29.5)
Hindu	125 (62.5)	92 (74.64)	33 (26.4)
Muslim	75 (37.5)	49 (60.34)	26 (34.66)

(Figures in line parenthesis show percentages.)

Out of 200 cases 125 (62.5%) cases were Hindus; out of which 74.64% cases were symptomatic and 26.4% were asymptomatic, while 75 (37.5%) were Muslims; out of which 60.34% were symptomatic and 34.66% were asymptomatic.

Table 5: Distribution of cases according to occupation and suggestive findings of RTI

Occupation	Total No. of cases 200	No. of patients	
		Symptomatic 141 (70.5)	Asymptomatic 59(29.5)
Housewives	139 (69.5)	91 (65.46)	48(34.54)
Working women	61 (30.5)	50 (29.50)	11(18.03)

(Figures in line parenthesis show percentages.)

Out of 200 cases 39 (69.5%) were housewives; out of which 65.46% were symptomatic and 34.54% were asymptomatic, while 61(30.5%) were working women; out of which 50(29.50%) were showing symptoms of RTI and 11(18.03%) were asymptomatic.

Table 6: Distribution of cases according to literacy and suggestive symptoms of RTI

Literacy	Total No. of cases 200	No. of patients	
		Symptomatic 141(70.5)	Asymptomatic 59(29.5)
illiterate	96 (48)	86 (90.90)	10 (10.4)
Literate	104 (52)	55 52.88	49 (47.2)

(Figures in line parenthesis show percentages.)

Out of 200 cases 96 (48%) were illiterate; out of which 86 (90.90%) were symptomatic for RTI and 10 (10.4%) were asymptomatic, while 104 (52%) cases were literate; out of which 55 (52.88%) were symptomatic and 49 (37.24%) were asymptomatic.

Table 7: Distribution of cases according to contraceptive used and suggestive findings of RTI

Contraception	Total No. of cases 200	No. of patients	
		Symptomatic 141(70.5)	Asymptomatic 59(29.5)
IUD	4 (2)	0	4(100)
Hormonal	12 (6)	1 (8.33)	11(91.67)
Tubectomy	88 (44)	76 (86.36)	12(13.64)
Male condom	30 (15)	4 (13.33)	26(86.67)
Non-users	66 (33)	60(90.90)	06(9.1)

(Figures in line parenthesis show percentages.)

Above table shows the current contraceptive practice in the study group. Out of 200 cases, 4 (2%) were IUD users, 12(6%) were hormonal pill users, 88(44%) had already undergone tubectomy and around 13.33% cases were dependant on their spouse for use of male condom as a contraceptive. None of the IUD users was found to be symptomatic, but among the hormonal pill users > 91% were asymptomatic. Those who resorted to permanent contraception almost 85% were symptomatic. Amongst the non-users the figure of symptomatic presentation was maximum almost 90%, while amongst those cases were dependent on their spouse for use of male condom as a contraceptive; the no. of symptomatic cases was around 14%.

Table 8: Symptom wise distribution of cases with respect to clinical diagnosis

Symptom	Bacterial Vaginosis	Trichomonas Vaginalis	Candidiasis
Vaginal Discharge	28 (43.75)	16 (40)	22 (59.75)
Pain in Abdomen	24 (37.5)	08 (20)	15 (90.54)
Backache	10 (15.62)	09 (22.5)	14 (37.83)
Itching	13 (20.31)	18 (45)	19 (51.35)
Increased Frequency of Micturition	15 (23.43)	11 (27.5)	11 (29.72)
Dyspareunia	07 (10.93)	05 (12.5)	09 (24.32)

(Figures in line parenthesis show percentages.)

Above table shows that out of 64 cases of Bacterial vaginosis, 43.75% complained of vaginal discharge, 37.5% complained of pain in abdomen, 15.62% patients complained of backache, 20.31% complained of itching at genital area, 23.43% complained of increased frequency of micturition and 10.93% complained of dyspareunia. Among 40 cases of Trichomonasvaginalis, vaginal discharge was present in 40% cases, pain in abdomen in 20%, backache in 22.5%, Itching in 45%, increased frequency of micturition in 27.5%, dyspareunia in 12.5% of cases. Among 36 cases of Candidiasis vaginal discharge was present in 59.45% cases, backache in 37.83%, pain in abdomen in 40.54%, itching at genital area in 51.35%, increased frequency of micturition in 29.72 % and dyspareunia in 24.32%.

DISCUSSION

Prevalence of RTI: Prevalence of RTI in our study is high than most of the other studies but less than the study of Aggarwal AK *et al*⁹.

Studies	Prevalence (%)
K.M. Bansal <i>et al</i> ⁵	40.50
SC Panda <i>et al</i> ⁶	39.2
Bhawana Pant <i>et al</i> ⁷	35
DeokiNandan <i>et al</i> ⁸	35.2
Aggarwal AK <i>et al</i> ⁹	70
Present study	63

As per the Different pathogens the prevalence was: The prevalence of Bacterial Vaginosis (32%) in our study is comparable to that in study done by Raya Gulati *et al*¹¹ (31%) and Oladele Teslim Ojuromi *et al*¹³ (34%) while it is more than study of Linda French *et al*¹² (21%) and Philippon *et al*¹⁴ (24%) but less than the study done by Danial V Landers *et al*¹⁰ (46%). Prevalence of Trichomonasvaginalis (11%) in the present study is comparable to study of Ray Gulati *et al*¹¹, while it is less than most of other studies. Prevalence of candidiasis was comparable to Study by Danial V Landers *et al*¹⁰ and Oladele Teslim Ojuromi *et al*¹³, while it was less than most of other studies. This difference among prevalence of infections over the world may be due to difference in geographic and socio-demographic factors. Knowing the approximate prevalence of different infections in their own setting may be helpful for physicians who have no laboratory facility.

Infection	Danial V Landers <i>et al</i> ¹⁰	Linda French <i>et al</i> ¹²	Ray A Gulati A.K. <i>et al</i> ¹¹	Oladele Teslim Ojuromi <i>et al</i> ¹³	Philippon <i>et al</i> ¹⁴	Present Study
Bacterial Vaginosis	46 %	21 %	31 %	34 %	24 %	32 %
TrichomonasVaginalis	29 %	8 %	11 %	30 %	14 %	11 %
Candidiasis	12 %	27 %	30 %	11 %	20 %	13.5%

Age Groups: Present study shows that the prevalence of RTI in age group ≤ 19 years and > 30 years is comparable to other studies, however the maximum prevalence was found in the age group of 21-29 years (56.44%) as that of other studies showing a relation with period of high fertility as well as it may be attributed to higher proportion of married people in the younger age which is sexually the most active age group.

Age Group (years)	K.M. Bansal <i>et al</i> ⁵ 200	Dasgupta Aprajita <i>et al</i> ¹⁵ 210	Bhawana Pant <i>et al</i> ⁷ 600	Present Study 126
≤ 19	57 (38.2)	68 (32)	219 (36.5)	35 (27.6)
20- 29	77 (51.52)	110 (52.43)	276 (46.0)	71 (56.44)
≥ 30	45 (30.5)	32 (15.23)	105 (17.5)	20 (16.00)

Parity: The prevalence of cases in nulliparous group in the present study is more than the other two studies, while for parity 3 - 4 and parity >4 it was less. These findings are comparable with S C Panda *et al*⁶ DeokiNandan *et al*⁸

Religion: The prevalence of cases of RTI in Muslims is almost comparable to other studies; however the prevalence of RTI cases in Muslims is higher as compared to Hindus this is comparable with K.M. Bansal⁵ who found 200 in Hindu 92 (46) Muslim 108 (54) and by Dasgupta *et al*¹⁴ it was 210; 84 (40) and 126 (60) and in Present Study 126; 53 (42) and 73 (58) respectively in Hindu and Muslims.

Occupation: Almost > 60% cases of RTI were housewives, which is higher than the working women. The prevalence of RTI cases was observed to be much less among the working women than their nonworking (housewife) counterparts (30% vs. 70%). This is because of their contribution to the family income, thereby empowering them to get involved in the decision making process in the family (e.g. use of condom, use of sanitary napkins, use of soap and water etc). This along with their higher literacy, better knowledge regarding menstrual hygiene, better and timely utilization of health care etc. go a long way in the maintenance of a better health status. These findings are comparable with Dasgupta *et al*¹⁴ (210) who found in Housewives 134 (64) and Working women 76 (36) Kantida *et al*¹⁶ (217) 143 (66) and 74 (34) and Present Study (126) 88 (70) and 38 (30) respectively in Housewives Working women.

Literacy: In the present study prevalence of RTI cases was found to be higher in illiterate (63.63%) as compared to literate (36.37%). This is a general phenomenon among the populations of Asia, including India. Substandard

hygiene, low socio-economic status, promiscuity and traditional taboos against openness about these diseases are the usual factors responsible for this high prevalence of RTI in illiterates. This is comparable with K.M. Bansal *et al*⁵ Dasgupta *et al*¹⁵.

Contraceptive Used: Prevalence of RTI according to the contraceptive practice in the present study population. The prevalence of RTI with contraceptive practice is different in various studies. The prevalence of RTI was highest in non-users (41.26%) as compared to other groups, which is comparable to other studies which also show high prevalence in non-users. It was lowest in the women who were dependent on their spouse for the use of condom as a contraceptive. These results of the study corroborate the universal fact that the use of condom during sexual intercourse prevents RTI.

Contraception	K.M. Bansal <i>et al</i> ⁵ 200	Dasgupta <i>et al</i> ¹⁵ 210	Present Study 126
IUD	30 (15)	25 (12)	04 (3.17)
Hormonal	7 (3.5)	10 (5)	09 (7.14)
Tubectomy	80 (40)	39 (17)	54 (43.65)
Male condom	15 (7.5)	23 (11)	6 (4.76)
Non-users	68 (34)	113 (55)	52 (41.26)

Symptom: The frequency of the symptoms with which cases presented. Cases presented with multiple complaints. There is variation in the frequency of the presenting symptoms of patients in our study group and other studies; however the most common symptom was vaginal discharge (44.5%) which is comparable to other studies, which also shows that it was the most common complaint. These findings are comparable with the following studies.

Symptom	Kantida <i>et al</i> ¹⁶ (%) 217	DeokiNandan <i>et al</i> ⁸ (%) 600	Present Study (%) 126
Vaginal discharge	79 (36)	252 (42)	81 (60)
Pain in Abdomen	53 (24.42)	120 (20)	30 (24)
Backache	-	132 (32)	50 (39)
Itching	42 (19.4)	90 (15)	27 (22)
Increased frequency of Micturition	-	120 (20)	19 (15)
Dyspareunia	33 (15.2)	60 (10)	25 (20)

CONCLUSION

It can be concluded from study that Majority of cases with suggestive symptoms of RTI were in the age group of 20-29 years & the asymptomatic cases were highest in the parity group > 4. Prevalence of RTI cases was high among Muslims nonworking (housewife), in illiterate and in non-users of contraceptives. In the study population cases were having multiple symptoms of which in overall vaginal discharge was the most common complaint.

REFERENCES

1. Kantida Chaijareenont, Korakot Sirimai, Dittakarn, Boriboonhirunsarn, Orawan Kiriwat. Accuracy of Nugent's Score and Each Amsel's Criteria in the Diagnosis of Bacterial Vaginosis. *J Med Assoc Thai* 2004; 87(11): 1270-4.
2. Aggarwal D. Reproductive tract infections – challenges and responses. *Health for the Millions*. 2001; 3:21–3.
3. Meheus AZ. Women's health and reproductive tract infections: The challenges posed by pelvic inflammatory disease, infertility, ectopic pregnancy and cervical cancer. In: Germain A, Holmes KK, Piot P, Wasserheit JN,

editors. Reproductive Tract Infections: Global impact and priorities for Women's Reproductive Health. New York: Plenum Press; 1992. pp. 61–91

4. National Guidelines on Prevention, Management and Control of Reproductive tract- Infections including Sexually Transmitted Infections. National AIDS Control Organization, MOHFW, Govt. of India; 2006. Nov.
5. Bansal KM, Singh K, Bhatangar S. Prevalence of lower RTI among married females in the reproductive age group (15-45). *Health Popul Perspect Issues* 2001; 24:157-63.
6. S C Panda, Lsarangi, D Bebartta, S Parida, O P Panigrahi. Prevalenc of RTI/STI among women of Reproductive Age in District Sundergarh (Orissa). *Indian Journal for the Practising* ;Vol.4, No.1(2007-03-2007-04)
7. Bhawna Pant, JV Singh, M Bhatnagar, SK Garg, H Chopra, SK Bajpai. Social Correlates in Reproductive Tract Infections among Married Women in Rural Area of Meerut. *Indian J Community Med.* 2008 January; 33(1): 52–53.
8. Deoki Nandan, S.K. Misra, AnitaSharma, Manish Jain. Estimation of Prevalenc of RTI/STD's among women of Reproductive Age Group in District Agra. *Indian Journal of Community Medicine*; Vol.27, No.3 (2002-07-2002-09).
9. Aggarwal AK, Kaur M, Kumar R. Community based study of reproductive tract infection among women of reproductive age in rural area of Haryana. *J Common Dis* 1993;31:223-8
10. Daniel V landers, Harold C Wiesenfeld, R Phillip Heine, Marijane A Krohn, Sharon L Hiller. Predictive value of the clinical diagnosis of lower genital tract infection in women. *American Journal of Obstetrics and Gynecology* vol 190, Issue 4, April 2004, Pages 1004-1008.
11. Ray A, Gulati AK, Pandey LK, Pandey S. Prevalence of common infective agents of vaginitis. *J Commun Dis* 1989 Sep; 21(3):241-4.
12. Linda French, Jennifer Horton, Michelle Matousek. Abnormal Vaginal discharge: Using office diagnostic testing more effectively. *The Journal of Family Practice*: October 2004. Vol 53, No. 10
13. Oladele TeslimOjoromi, Wellington Aghoghowia Oyibo, Adetokunbo OlufelaTayo, Comfort Aeejokelbidapo, Adetayo Foluscho Fagbenro-Beyioku, Oladipo Olarinre Oladosu, I.Isiaka Olaniyi Ola-Gbadamosi, Eromese Ruth Okposugbo, and Abiola Oluwatoyin Balogun. Reliance on microscopy in T. vaginalis diagnosis & its prevalence in females presenting with vaginal discharge in lagos, Nigeria. *J infect Developing Countries* 2007; 1 (2): 210-213
14. B BLÔ, M Philippon, P Cunin, D Meynard, M Tandia-igina. The microbial etiology of genital discharges in Nouakchott, Mauritania. *Bulletin de la Société de pathologie exotique* (1990): Volume: 90 ISSN: 0037-9085 ISO Abbreviation: Bull SocPatholExot Publication Date: 1997.
15. Dasgupta Aprajita, Sarkar Madhutandra. A study on reproductive tract infections among married women in the reproductive age group (15-45 yrs) in a slum of Kolkata. *J ObstetGynecol India* November/December 2008; Vol.58, No.6: pg 518-522.
16. Kantida Chaijareenont, Korakot Sirimai, Dittakarn, Boriboonhirunsarn, Orawan Kiriwat. Accuracy of Nugent's Score and Each Amsel's Criteria in the Diagnosis of Bacterial Vaginosis. *J Med Assoc Thai* 2004; 87(11): 1270-4.

Source of Support: None Declared
Conflict of Interest: None Declared