

Almost sure limit points of maxima of stationary gaussian sequences through at least geometrically fast subsequences

S S Nayak^{1*}, Varalaxmi T Shedole²

¹Department of Statistics, Gulbarga University, Gulbarga-585106, Karnataka, INDIA.

²Government first Grade College, Hubli, Karnataka, INDIA.

Email: ssnayak2006@rediffmail.com

Abstract

Let $(X_n, n \geq 1)$ be a discrete - parameter Stationary Gaussian process with $E(X_i)=0, EX_i^2=1$ for all i and $E(X_i X_{i+n})=r(n)$. Let $(Y_n, n \geq 1)$ be an independent copy of $(X_n, n \geq 1)$. Let $M_{1n} = \max(X_1, X_2, \dots, X_n), M_{2n} = \max(Y_1, Y_2, \dots, Y_n), U_n = \frac{(M_{1n}-b_n)}{a_n}$ and $V_n = \frac{(M_{2n}-b_n)}{a_n}$ where $b_n = (2 \log n)^{1/2}$ and $a_n = (\log \log n)(2 \log n)^{-1/2}$. Let (n_k) be a subsequence of positive integers that is at least geometrically fast. Under the condition that either $(\log n)^{1+\gamma}r(n) = O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$ or $\sum_{j=1}^{\infty} r^2(j) < \infty$ the set of all almost sure limit points of the vector sequence $\left\{ U_n = \frac{(M_{1n}-b_n)}{a_n}, V_n = \frac{(M_{2n}-b_n)}{a_n} \right\}$ is obtained.

Keywords: stationary gaussian sequence.

*Address for Correspondence:

Dr. S. S. Nayak, Department of Statistics, Gulbarga University, Gulbarga-585106, Karnataka, INDIA.

Email: chhayajadhav@gmail.com

Received Date: 27/02/2015 Accepted Date: 11/03/2015

Access this article online

Quick Response Code:



Website:

www.statperson.com

DOI: 12 March 2015

INTRODUCTION

Let $(X_n, n \geq 1)$ be a discrete - parameter Stationary Gaussian process with $E(X_i)=0, EX_i^2=1$ for all i and $E(X_i X_{i+n})=r(n)$. Let $(Y_n, n \geq 1)$ be an independent copy of $(X_n, n \geq 1)$. Let $M_{1n} = \max(X_1, X_2, \dots, X_n), M_{2n} = \max(Y_1, Y_2, \dots, Y_n), U_n = \frac{(M_{1n}-b_n)}{a_n}$ and $V_n = \frac{(M_{2n}-b_n)}{a_n}$ where $b_n = (2 \log n)^{1/2}$ and $a_n = (\log \log n)(2 \log n)^{-1/2}$. Pickands (1969) established that if either $(\log n)^\alpha r(n) \rightarrow 0$ as $n \rightarrow \infty$ for some $\alpha > 0$ or $\sum_{j=1}^{\infty} r^2(j) < \infty$ then almost surely, as $n \rightarrow \infty$ $\limsup U_n = \frac{1}{2}$ and $\liminf U_n = -\frac{1}{2}$. Mittal (1974) showed that the above results continue to hold if the condition $(\log n)^\alpha r(n) \rightarrow 0$ as $n \rightarrow \infty$ for some $\alpha > 0$ is replaced by the weaker condition $(\log n)^\alpha r(n) = O(1)$ as $n \rightarrow \infty$ for some $\alpha > 0$. Vishnu Hebbar (1980) obtained the almost sure limit set of (U_n, V_n) when either $(\log n)^{1+\gamma}r(n) = O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$ or $\sum_{j=1}^{\infty} r^2(j) < \infty$. Let (n_k) be a subsequence of positive integers with $n_k \rightarrow \infty$ as $k \rightarrow \infty$. It is said to be at least geometrically fast if $\limsup \left(\frac{n_k}{n_{k+1}} \right) < 1$. Vasudeva and Savitha (1995) established the law of the iterated logarithm for (U_{n_k}) . In this paper, we

extend their result to the vector case by finding the almost sure limit set of (U_{n_k}, V_{n_k}) . Almost sure limit points of random vectors with independent components have received considerable attention in literature. One can refer the papers by LePage (1973), Pakshirajan and Vasudeva (1977), Strassen (1964), Nayak,S.S. (1984, 1985, 1986, 1988,1994,2000 and 2001) and the references therein. Throughout the paper, const. stands for a positive constant which may have different values at different appearances. “infinitely often” is written as i.o.

PRELIMINARIES

Let (n_k) be a subsequence of positive integers such that $n_k \rightarrow \infty$ as $k \rightarrow \infty$ and $\limsup \left(\frac{n_k}{n_{k+1}}\right) < 1$. Let $\varepsilon^* = \inf\{\varepsilon: \sum(\log n_k)^{-(\varepsilon+1/2)} < \infty\}$. For $-\frac{1}{2} < x_i \leq \varepsilon^*$, $i=1,2$ with $-1 < x_1+x_2 \leq \varepsilon^* - \frac{1}{2}$, let $n_k^* = n_{u(k)}$ where $u(k) = \left\lceil k^{\frac{1}{1+x_1+x_2}} \right\rceil$ and $[x]$ is the greatest integer $\leq x$. Let $m_k = \left\lceil n_k^* (\log k)^{-\frac{1}{2}} \right\rceil$. Let us define the events $F_k = \{\max_{n_k^* - m_k + 1 \leq j \leq n_k^*} X_j > d_{n_k^*}(x_1)\}$ and $G_k = \{\max_{n_k^* - m_k + 1 \leq j \leq n_k^*} Y_j > d_{n_k^*}(x_2)\}$ where $d_{n_k^*}(x) = a_{n_k^*} x + b_{n_k^*}$. When $r(n)=0$, the corresponding events are respectively denoted by F_k^* and G_k^* . Lemma 2.1 (Vasudeva and Savitha, 1995): Assume that either $(\log n)^{1+\gamma}r(n)=O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$ or $\sum_{j=1}^{\infty} r^2(j) < \infty$. Then $\limsup U_{n_k} = \varepsilon^*$ a.s. Lemma 2.2 (Vasudeva and Savitha, 1995): Assume that either $(\log n)^{1+\gamma}r(n)=O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$ or $\sum_{j=1}^{\infty} r^2(j) < \infty$. Then for any subsequence (v_k) of positive integers with $\lim_{k \rightarrow \infty} v_k = \infty$ we have $\liminf U_{v_k} = -\frac{1}{2}$. Lemma 2.3: Let $0 < \alpha < \frac{1-\delta}{1+\delta}$ and

$$A_1(k) = \sum_{j=1}^{\lfloor m_k^\alpha \rfloor} |r(j)| (m_k - j)(1 - r^2(j))^{-1/2} \exp\left\{-\frac{d_{n_k^*}^2(x_1)}{(1+|r(j)|)}\right\}$$

where $\sup_{n \geq 1} |r(n)| = \delta$ ($0 < \delta < 1$).

Then $\sum_{k=1}^{\infty} P(G_k^*)A_1(k) < \infty$.

Proof: Stationarity implies $\sup_{n \geq 1} |r(n)| = \delta$ ($0 < \delta < 1$). We have

$$\begin{aligned} A_1(k) &< (\text{const.})(n_k^*)^{-\frac{2}{1+\delta}} (\log n_k^*)^{-\frac{2x_1}{1+\delta}} \sum_{j=1}^{\lfloor m_k^\alpha \rfloor} (m_k - j) \exp\left\{-\frac{1}{1+\delta} (2 \log n_k^* + 2x_1 \log \log n_k^*) + o(1)\right\} \\ &< (\text{const.}) m_k^{\alpha+1} (n_k^*)^{-\frac{2}{1+\delta}} (\log n_k^*)^{-\frac{2x_1}{1+\delta}}, k \geq k_1 \\ &< (\text{const.})(\log k)^{\frac{\alpha+1}{2}} (\log n_k^*)^{-\frac{2x_1}{1+\delta}} (n_k^*)^{\alpha+1-\frac{2}{1+\delta}}, k \geq k_1. \end{aligned}$$

Also, $P(G_k^*) = 1 - \Phi^{m_k} \left(d_{n_k^*}(x_2) \right)$ where Φ is the d.f. of a standard normal random variable.

$$\begin{aligned} &= 1 - \exp\left\{m_k \log \left\{1 - \left(1 - \Phi \left(d_{n_k^*}(x_2)\right)\right)\right\}\right\} \\ &= 1 - \exp\left\{-m_k \left(1 - \Phi \left(d_{n_k^*}(x_2)\right)\right) (1 + o(1))\right\} \\ &< (\text{const.})(\log k)^{-1/2} (\log n_k^*)^{-(x_2+1/2)} k \geq k_2 \text{ by the tail behavior of the standard normal distribution.} \end{aligned}$$

Let $N = \max(k_1, k_2)$. Then

$$\sum_{k=N}^{\infty} P(G_k^*)A_1(k) < (\text{const.}) \sum_{k=N}^{\infty} (\log k)^{-\frac{\alpha+2}{2}} (n_k^*)^{-\theta} (\log n_k^*)^{-\beta}$$

where $\theta = \frac{2}{1+\delta} - \alpha - 1 > 0$

$$\text{and } \beta = \frac{2x_1}{1+\delta} + x_2 + \frac{1}{2}.$$

$$\begin{aligned} &< (\text{const.}) \sum_{k=N}^{\infty} (n_k^*)^{-\theta} (\log n_k^*)^{-\beta} \\ &= (\text{const.}) \sum_{k=N}^{\infty} e^{-\theta \log n_k^*} (\log n_k^*)^{-\beta} \\ &< (\text{const.}) \sum_{k=N}^{\infty} (\log n_k^*)^{-(l+\beta)} \text{ where } l > \max(1, 1 + x_1 + x_2 - \beta) \text{ is an integer.} \end{aligned}$$

Since $\limsup \frac{n_k}{n_{k+1}} < 1$, we have $n_k^* > (\text{const.})a^{u(k)}$ for $k \geq k_3$ where $a > 1$ and $u(k) = \left\lceil k^{\frac{1}{1+x_1+x_2}} \right\rceil$. Hence $\sum_{k=N}^{\infty} P(G_k^*)A_1(k) <$

$$(\text{const.}) + (\text{const.}) \sum_{k=k_3}^{\infty} k^{-\frac{l+\beta}{1+x_1+x_2}} < \infty.$$

Lemma 2.4: Let $0 < \alpha < \frac{1-\delta}{1+\delta}$ where $\sup_{n \geq 1} |r(n)| = \delta$ ($0 < \delta < 1$). Let $\sup_{k \geq n} |r(k)| = \delta(n)$ and

$$\begin{aligned} A_2(k) &= \sum_{j=\lfloor m_k^\alpha \rfloor + 1}^{m_k - 1} |r(j)| (m_k - j)(1 - r^2(j))^{-1/2} \exp\left\{-\frac{d_{n_k^*}^2(x_1)}{(1+|r(j)|)}\right\}. \text{ Assume that either} \\ &(\log n)^{1+\gamma}r(n)=O(1) \text{ as } n \rightarrow \infty \text{ for some } \gamma > 0 \text{ or } \sum_{j=1}^{\infty} r^2(j) < \infty. \text{ Then } \sum_{k=1}^{\infty} P(G_k^*)A_2(k) < \infty. \end{aligned}$$

Proof: First let $(\log n)^{1+\gamma}r(n)=O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$. Then $(\log n)^{1+\gamma} \delta(n)=O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$. Now

$$A_2(k) < (\text{const.})\delta(m_k^\alpha)(1 - \delta^2)^{-1/2}(n_k^*)^{-\frac{2}{1+\delta(m_k^\alpha)}} (\log n_k^*)^{-\frac{2}{1+\delta(m_k^\alpha)}} \sum_{j=[m_k^\alpha]+1}^{m_k-1} (m_k-j), k \geq k_4$$

$$< (\text{const.}) \delta(m_k^\alpha) m_k^2 (n_k^*)^{-\frac{2}{1+\delta(m_k^\alpha)}} (\log n_k^*)^{-\frac{2x_1}{1+\delta(m_k^\alpha)}}, k \geq k_5$$

$$< (\text{const.}) \delta(m_k^\alpha) m_k^{2\delta(m_k^\alpha)} (\log k)^{\delta(m_k^\alpha)} (\log n_k^*)^{-\frac{2x_1}{1+\delta(m_k^\alpha)}}, k \geq k_6 \text{ since } m_k \sim n_k^* (\log k)^{-1/2} \text{ as } k \rightarrow \infty.$$

$$=(\text{const.}) \delta(m_k^\alpha) (\log k)^{\delta(m_k^\alpha)} (\log n_k^*)^{-\frac{2x_1}{1+\delta(m_k^\alpha)}} \exp\{2\delta(m_k^\alpha) \log m_k\}, k \geq k_6$$

$$< (\text{const.}) (\log m_k)^{-\gamma-1} (\log k)^{\delta(m_k^\alpha)} (\log n_k^*)^{-\frac{2x_1}{1+\delta(m_k^\alpha)}}, k \geq k_7 \text{ since } \delta(m_k^\alpha) (\log m_k^\alpha)^{1+\gamma} = O(1) \text{ as } k \rightarrow \infty.$$

Since $\limsup \frac{n_k}{n_{k+1}} < 1$, we have $n_k^* > (\text{const.})a^{u(k)}$ for k large where $a > 1$ and $u(k) = \left\lceil k^{\frac{1}{1+x_1+x_2}} \right\rceil$. This implies that $\frac{\log \log k}{\log n_k^*}$

$\rightarrow 0$ as $k \rightarrow \infty$. Hence $\log m_k \sim \log n_k^*$ as $k \rightarrow \infty$. Hence

$$A_2(k) < (\text{const.}) (\log k)^{\delta(m_k^\alpha)} (\log n_k^*)^{-\frac{2x_1}{1+\delta(m_k^\alpha)}-\gamma-1}, k \geq k_8.$$

Also, $P(G_k^*) < (\text{const.}) (\log k)^{-1/2} (\log n_k^*)^{-(x_2+1/2)}$ for large k by the tail behavior of the standard normal distribution. Hence

$$P(G_k^*)A_2(k) < (\text{const.}) (\log k)^{\delta(m_k^\alpha)-1/2} (\log n_k^*)^{-\frac{2x_1}{1+\delta(m_k^\alpha)}-x_2-\gamma-3/2}, k \geq k_9.$$

$$< (\text{const.}) (\log k)^{\delta(m_k^\alpha)-1/2} k^{-\left(\frac{2x_1}{1+\delta(m_k^\alpha)}+x_2+\gamma+3/2\right)/(1+x_1+x_2)}, k \geq k_{10} \text{ since } \limsup \frac{n_k}{n_{k+1}} < 1 \text{ implies}$$

$$n_k^* > (\text{const.})a^{u(k)} \text{ for } k \text{ large where } a > 1 \text{ and } u(k) = \left\lceil k^{\frac{1}{1+x_1+x_2}} \right\rceil.$$

Note that $(\log n)^{1+\gamma} \delta(n) = O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$ implies $\delta(m_k^\alpha) \rightarrow 0$ as $k \rightarrow \infty$.

Let $0 < \epsilon_1 < \frac{x_1+\gamma+1/2}{1+x_1+x_2}$. This is possible since $x_1 > -\frac{1}{2}$ and $1+x_1+x_2 > 0$. Hence

$$\sum_{k=1}^{\infty} P(G_k^*)A_2(k) < \text{const.} + (\text{const.}) \sum_{k=k_{11}}^{\infty} (\log k)^{\epsilon_1-1/2} k^{\left(\epsilon_1 - \frac{2x_1+x_2+\gamma+3/2}{1+x_1+x_2}\right)} < \infty.$$

Now let $\sum_{j=1}^{\infty} r^2(j) < \infty$. By Cauchy-Schwarz inequality we have

$$(A_2(k))^2 < \sum_{j=[m_k^\alpha]+1}^{m_k-1} r^2(j) \sum_{j=[m_k^\alpha]+1}^{m_k-1} (m_k-j)^2 (1-r^2(j)) \times$$

$$\exp\left\{-\frac{2}{1+|r(j)|} \left\{2 \log n_k^* + 2x_1 \log \log n_k^* + x_1^2 \frac{(\log \log n_k^*)^2}{2 \log \log n_k^*}\right\}\right\}$$

$$< (\text{const.}) \left\{ \sum_{j=1}^{\infty} r^2(j) \right\} (n_k^*)^{-\frac{4}{1+\delta(m_k^\alpha)}} (\log n_k^*)^{-\frac{4x_1}{1+\delta(m_k^\alpha)}} \sum_{j=[m_k^\alpha]+1}^{m_k-1} (m_k-j)^2, k \geq k_{11}$$

$$< (\text{const.}) \left\{ \sum_{j=1}^{\infty} r^2(j) \right\} (n_k^*)^{-\frac{4}{1+\delta(m_k^\alpha)}} (\log n_k^*)^{-\frac{4x_1}{1+\delta(m_k^\alpha)}} m_k^3, k \geq k_{12}.$$

$$< (\text{const.}) \left\{ \sum_{j=1}^{\infty} r^2(j) \right\} (n_k^*)^{3-\frac{4}{1+\delta(m_k^\alpha)}} (\log k)^{-3/2} (\log n_k^*)^{-\frac{4x_1}{1+\delta(m_k^\alpha)}}, k \geq k_{13} \text{ since } m_k = \left\lceil n_k^* (\log k)^{-\frac{1}{2}} \right\rceil.$$

$$\text{Hence } A_2(k) < (\text{const.}) \left\{ \sum_{j=1}^{\infty} r^2(j) \right\}^{1/2} (n_k^*)^{3/2-\frac{2}{1+\delta(m_k^\alpha)}} (\log k)^{-3/4} (\log n_k^*)^{-\frac{2x_1}{1+\delta(m_k^\alpha)}}, k \geq k_{13}$$

$$< (\text{const.}) \left\{ \sum_{j=1}^{\infty} r^2(j) \right\}^{1/2} (n_k^*)^{\epsilon_1-1/2} (\log k)^{-3/4} (\log n_k^*)^{\epsilon_1-2x_1}, k \geq k_{14}, 0 < \epsilon_1 < \frac{1}{2},$$

since $\delta(m_k^\alpha) \rightarrow 0$ as $k \rightarrow \infty$.

$$< (\text{const.}) \left\{ \sum_{j=1}^{\infty} r^2(j) \right\}^{1/2} (\log n_k^*)^{\epsilon_1-2x_1} \exp\{-\theta \log n_k^*\}, k \geq k_{15} \text{ where } \theta = \frac{1}{2} - \epsilon_1.$$

$$< (\text{const.}) \left\{ \sum_{j=1}^{\infty} r^2(j) \right\}^{1/2} (\log n_k^*)^{\epsilon_1-2x_1-l}, k \geq k_{15} \text{ where } l \text{ is an integer such that}$$

$$l \geq \max(1, \epsilon_1^* + \epsilon_1 - 2x_1 - x_2).$$

Also $P(G_k^*) < (\text{const.}) (\log k)^{-1/2} (\log n_k^*)^{-(x_2+1/2)}$ for large k by the tail behavior of the standard normal distribution. Hence,

Hence,

$$\sum_{k=1}^{\infty} P(G_k^*)A_2(k) < (\text{const.}) + (\text{const.}) \left\{ \sum_{j=1}^{\infty} r^2(j) \right\}^{1/2} \sum_{k=k_{16}}^{\infty} (\log n_k^*)^{\epsilon_1-2x_1-l-x_2-1/2} < \infty.$$

Lemma 2.5: Let I_k be the indicator function of the event E_k . Assume that $(\log n)^{1+\gamma}r(n)=O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$.

Then $\sup_n |\sum_{N \leq k < l \leq N} \text{Cov}(I_k, I_l)| < \infty$ where N is a large positive integer.

Proof: We have for $k < l$,

$$\begin{aligned} Cov(I_k, I_l) &= E(I_k I_l) - E(I_k) E(I_l) \\ &= P(F_k \cap F_l) P(G_k \cap G_l) - P(F_k) P(F_l) P(G_k) P(G_l) \\ &= \{P(F_k \cap F_l) - P(F_k)P(F_l)\} \{P(G_k \cap G_l) - P(G_k)P(G_l)\} + P(G_k)P(G_l) \\ &\quad + P(F_k) P(F_l) \{P(G_k \cap G_l) - P(G_k)P(G_l)\}. \text{ Therefore} \\ |Cov(I_k, I_l)| &\leq \{P(F_k \cap F_l) - P(F_k)P(F_l)\} \{P(G_k \cap G_l) - P(G_k)P(G_l)\} \\ &\quad + P(G_k)P(G_l) \{P(F_k \cap F_l) - P(F_k)P(F_l)\} + P(F_k) P(F_l) \{P(G_k \cap G_l) - P(G_k)P(G_l)\}. \dots \end{aligned} \tag{2.1}$$

By Qualls and Watanabe (1971) we get

$$|P(F_k \cap F_l) - P(F_k)P(F_l)| = |P(F_k^c \cap F_l^c) - P(F_k^c)P(F_l^c)| \tag{2.2}$$

$\leq \sum_{\mu=1}^{m_l} \sum_{v=1}^{m_k} |r| \int_0^1 \varphi(d_{n_k^*}(x_1), d_{n_l^*}(x_1); \lambda r) d\lambda$ where $\varphi(u, v, \rho)$ is the standard bivariate normal density with correlation coefficient ρ and $r = r(n_k^* - m_l + \mu - n_k^* + m_k - v)$. Since $(\log n)^{1+\gamma} r(n) = O(1)$ as $n \rightarrow \infty$, we have

$(\log n)^{1+\gamma} \delta(n) = O(1)$ as $n \rightarrow \infty$ where $\sup_{k \geq n} |r(k)| = \delta(n)$. Stationarity of X_n 's and the condition on $r(n)$ ensure that

$\delta(1) < 1$. Let $\limsup \frac{n_k^*}{n_{k+1}^*} = a$ ($0 < a < 1$). Hence, noting that $\frac{m_l}{n_l^*} \rightarrow 0$ as $l \rightarrow \infty$, we have, $n_l^* - m_l + \mu - n_k^* + m_k - v \geq (const.) n_l^*$ for l large. Therefore $|n_l^* - m_l + \mu - n_k^* + m_k - v| \leq \delta((const.) n_l^*) \leq (const.) (\log n_l^*)^{-1-\gamma}$ for large k and l such that $l > k \geq N$, N being a sufficiently large positive integer.

Note that

$$\begin{aligned} \varphi(d_{n_k^*}(x_1), d_{n_l^*}(x_1); \lambda r) &\leq (2\pi)^{-1} \left((1 - \delta(1)) \right)^{-1/2} \times \\ \exp\left\{-\frac{1}{2} \left(d_{n_k^*}^2(x_1) - 2|r| d_{n_k^*}(x_1) d_{n_l^*}(x_1) + d_{n_l^*}^2(x_1) \right)\right\} \end{aligned} \tag{2.3}$$

$\leq (const.) \exp\left\{-\frac{1}{2} \left(d_{n_k^*}^2(x_1) + (1 - 2|r|) d_{n_l^*}^2(x_1) \right)\right\}$ because $d_{n_j^*}$'s are monotonically increasing in j .

$< (const.) (n_k^*)^{-1} (\log n_k^*)^{-x_1} (n_l^* (\log n_l^*)^{x_1})^{-(1-2\delta((const.) n_l^*))}$
 $= (const.) (n_k^*)^{-1} (\log n_k^*)^{-x_1} (n_l^*)^{-1} (\log n_l^*)^{-x_1} \exp\{2\delta((const.) n_l^*) (\log n_l^* + x_1 \log \log n_l^*)\}$
 $< (const.) (n_k^*)^{-1} (\log n_k^*)^{-x_1} (n_l^*)^{-1} (\log n_l^*)^{-x_1} \exp\{2(const.) (\log n_l^*)^{-1-\gamma} (\log n_l^* + x_1 \log \log n_l^*)\}$
 since $(\log n)^{1+\gamma} \delta(n) = O(1)$ as $n \rightarrow \infty$.

$< (const.) (n_k^*)^{-1} (\log n_k^*)^{-x_1} (n_l^*)^{-1} (\log n_l^*)^{-x_1}$.

Hence the R.H.S of (2.2) can be majorized by

$$\begin{aligned} (const.) m_k m_l \delta((const.) n_l^*) (n_k^*)^{-1} (\log n_k^*)^{-x_1} (n_l^*)^{-1} (\log n_l^*)^{-x_1} \\ \leq (const.) (\log k)^{-1/2} (\log l)^{-1/2} (\log n_k^*)^{-x_1} (\log n_l^*)^{-x_1-\gamma-1} \end{aligned} \tag{2.4}$$

Similarly $|P(G_k \cap G_l) - P(G_k)P(G_l)|$ can be majorized by an expression which is obtained from (2.4)

By replacing x_1 by x_2 . Hence the first term of R.H.S of (2.1) is

$$\begin{aligned} \leq (const.) (\log k)^{-1} (\log l)^{-1} (\log n_k^*)^{-(x_1+x_2)} (\log n_l^*)^{-(x_1+x_2+2\gamma+2)} \\ \leq (const.) (\log k)^{-1} (\log n_k^*)^{-(x_1+x_2+\gamma+1)} (\log l)^{-1} (\log n_l^*)^{-(x_1+x_2+\gamma+1)}. \end{aligned}$$

The second term of (2.1) for large $l > k \geq N$ large is

$$\begin{aligned} \leq m_k P(Y_1 > d_{n_k^*}(x_2)) m_l P(Y_1 > d_{n_l^*}(x_2)) \{P(F_k \cap F_l) - P(F_k)P(F_l)\} \\ \leq (const.) m_k m_l (\log k)^{-3/2} (\log l)^{-1} (\log n_k^*)^{-(x_1+x_2+1/2)} (\log n_l^*)^{-(x_1+x_2+\gamma+3/2)} \\ \leq (const.) (\log k)^{-1} (\log n_k^*)^{-(x_1+x_2-1/2)} (\log n_l^*)^{-(x_1+x_2+\gamma+3/2)} \\ \leq (const.) (\log k)^{-1} (\log n_k^*)^{-(x_1+x_2+\gamma/2+1)} (\log l)^{-1} (\log n_l^*)^{-(x_1+x_2+\gamma/2+1)}. \end{aligned}$$

The third term of (2.1) is bounded by the same expression as the second. From these bounds the proof the lemma is complete.

Lemma 2.6: Let I_k be the indicator function of the event E_k . Assume that $\sum_{j=1}^{\infty} r^2(j) < \infty$. Then

$\sup_n | \sum_{N \leq k < l \leq n} Cov(I_k, I_l) | < \infty$ where N is a large positive integer.

Proof: We have

$$\begin{aligned} |Cov(I_k, I_l)| &\leq |P(G_k \cap G_l) - P(G_k)P(G_l)| + |P(F_k \cap F_l) - P(F_k)P(F_l)| \\ &\leq \sum_{\mu=1}^{m_l} \sum_{v=1}^{m_k} |r| \left(\int_0^1 \varphi(d_{n_k^*}(x_1), d_{n_l^*}(x_1); \lambda r) d\lambda + \int_0^1 \varphi(d_{n_k^*}(x_2), d_{n_l^*}(x_2); \lambda r) d\lambda \right) \end{aligned}$$

by lemma 1.5 of Qualls and Watanabe(1971) where r is defined at (2.2) of the proof of lemma 2.5.

The above expression can be majorized by

$$\sum_{\mu=1}^{m_l} \sum_{v=1}^{m_k} |r| \left(\exp \left\{ -\frac{1}{2} \left(d_{n_k^*}^2(x_1) + (1 - 2|r|)d_{n_l^*}^2(x_1) \right) \right\} + \exp \left\{ -\frac{1}{2} \left(d_{n_k^*}^2(x_2) + (1 - 2|r|)d_{n_l^*}^2(x_2) \right) \right\} \right) \quad (2.5)$$

(c.f. (2.3) of the proof of lemma 2.5)

As in (2.2) of the proof of lemma 2.5, we have $n_l^* - m_l + \mu - n_k^* + m_k - v \geq (\text{const.})n_l^*$ for l large.

Since $r(n) \rightarrow 0$ as $n \rightarrow \infty$, we get

$|r| = |r(n_l^* - m_l + \mu - n_k^* + m_k - v)| < \varepsilon$ for $l > k \geq N$ where $\varepsilon > 0$ is sufficiently small. Hence (2.5) can be majorized by

$$\left(\exp \left\{ -\frac{1}{2} \left(d_{n_k^*}^2(x_1) + (1 - 2\varepsilon)d_{n_l^*}^2(x_1) \right) \right\} + \exp \left\{ -\frac{1}{2} \left(d_{n_k^*}^2(x_2) + (1 - 2\varepsilon)d_{n_l^*}^2(x_2) \right) \right\} \right) \times \sum_{\mu=1}^{m_l} \sum_{v=1}^{m_k} |r(n_l^* - m_l + \mu - n_k^* + m_k - v)| \quad (2.6)$$

By Cauchy-Schwarz inequality

$$\begin{aligned} & \sum_{\mu=1}^{m_l} \sum_{v=1}^{m_k} |r(n_l^* - m_l + \mu - n_k^* + m_k - v)| \\ & \leq m_l^{1/2} \left(\sum_{\mu=1}^{m_l} \left(\sum_{v=1}^{m_k} |r(n_l^* - m_l + \mu - n_k^* + m_k - v)| \right)^2 \right)^{1/2} \\ & \leq m_l^{1/2} m_k^{1/2} \left(\sum_{\mu=1}^{m_l} \sum_{v=1}^{m_k} r^2 (n_l^* - m_l + \mu - n_k^* + m_k - v) \right)^{1/2} \\ & \leq m_l^2 m_k \left(\sum_{j=1}^{\infty} r^2(j) \right)^{1/2} \text{ since} \\ & \sum_{\mu=1}^{m_l} \sum_{v=1}^{m_k} r^2 (n_l^* - m_l + \mu - n_k^* + m_k - v) \\ & = \sum_{\mu=1}^{m_l} (r^2(n_l^* - m_l + \mu - n_k^* + m_k - 1) + \dots + r^2(n_l^* - m_l + \mu - n_k^*)) \\ & \leq m_k \left(\sum_{j=1}^{\infty} r^2(j) \right)^{1/2}. \end{aligned}$$

Thus (2.6) is bounded by

$$(\text{const.}) m_l^2 m_k (n_k^*)^{-1} (n_l^*)^{-(1-2\varepsilon)} \left((\log n_k^*)^{-x_2} (\log n_l^*)^{-(1-2\varepsilon)x_2} + (\log n_k^*)^{-x_1} (\log n_l^*)^{-(1-2\varepsilon)x_1} \right)$$

This complete the proof of the lemma.

THE MAIN RESULT

Theorem 3.1: Assume that either $(\log n)^{1+\gamma}r(n)=O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$ or $\sum_{j=1}^{\infty} r^2(j) < \infty$. Let (n_k) be any subsequence of positive integers such that $n_k \rightarrow \infty$ as $k \rightarrow \infty$ and $\limsup \frac{n_k}{n_{k+1}} < 1$. Then the set of all almost sure limit points of (U_{n_k}, V_{n_k}) is

$$S_2 = \left\{ (x_1, x_2) : -\frac{1}{2} \leq x_1, x_2 \leq \frac{1}{2}, x_1 + x_2 \leq \varepsilon^* - \frac{1}{2} \right\} \text{ where } \varepsilon^* = \inf \left\{ \varepsilon : \sum (\log n_k)^{-(\varepsilon+1/2)} < \infty \right\}.$$

The proof of the theorem is based on the following three lemmas. Let n_k^*, m_k, F_k and G_k be as in section 2.

Lemma 3.1: For all $x_1, x_2 > -\frac{1}{2}$ and for all $\varepsilon > 0$, we have

$$(i) P(U_{n_k^*} > x_1 + \varepsilon, V_{n_k^*} > x_2 \text{ i. o.}) = 0 \text{ and } (ii) P(U_{n_k^*} > x_1, V_{n_k^*} > x_2 + \varepsilon \text{ i. o.}) = 0$$

Proof: Since $U_{n_k^*}$ and $V_{n_k^*}$ are independent,

$$P(U_{n_k^*} > x_1 + \varepsilon, V_{n_k^*} > x_2) = P(U_{n_k^*} > x_1 + \varepsilon) P(V_{n_k^*} > x_2)$$

$$\leq n_k^{*2} P(X_1 > d_{n_k^*}(x_1 + \varepsilon)) P(Y_1 > d_{n_k^*}(x_2))$$

$\sim (\text{const.}) (\log n_k^*)^{-(1+x_1+x_2+\varepsilon)}$ as $k \rightarrow \infty$, using the known result

$1 - \Phi(x) \sim (2\pi)^{-1/2} x^{-1} \exp\left(-\frac{x^2}{2}\right)$ as $x \rightarrow \infty$ for the standard normal distribution function Φ . Since $\limsup \frac{n_k}{n_{k+1}} < 1$, we

have $n_k > a^k, 0 < a < 1$ for k large. Hence $\log n_k^* > (\text{const.})u(k)$ for $k \geq k_0$ where $u(k) = \left\lfloor k^{\frac{1}{1+x_1+x_2}} \right\rfloor$ and $[x]$ is the greatest

integer $\leq x$. Thus $\sum_k P(U_{n_k^*} > x_1 + \varepsilon, V_{n_k^*} > x_2) < \infty$. An application of the Borel-Cantelli lemma completes the proof of (i). Proof of (ii) is similar.

Lemma 3.2: Assume that either $(\log n)^{1+\gamma}r(n)=O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$ or $\sum_{j=1}^{\infty} r^2(j) < \infty$.

For all $x_1, x_2 > -\frac{1}{2}$ with $x_1 + x_2 \leq \varepsilon^* - \frac{1}{2}$, we have

$$P(U_{n_k^*} > x_1, V_{n_k^*} > x_2 \text{ i. o.}) = 1.$$

Proof: Recall that let $n_k^* = n_{u(k)}$ where $u(k) = \left\lceil k^{\frac{1}{1+x_1+x_2}} \right\rceil$ and $[x]$ is the greatest integer $\leq x$, $m_k = \left\lceil n_k^* (\log k)^{-\frac{1}{2}} \right\rceil$, F_k

$$= \{ \max_{n_k^* - m_k + 1 \leq j \leq n_k^*} X_j > d_{n_k^*}(x_1) \} \text{ and,}$$

$G_k = \{ \max_{n_k^* - m_k + 1 \leq j \leq n_k^*} Y_j > d_{n_k^*}(x_2) \}$ where $d_{n_k^*}(x) = a_{n_k^*}x + b_{n_k^*}$. When $r(n) = 0$, the corresponding events are respectively denoted by F_k^* and G_k^* . Define $E_k = F_k \cap G_k$. Observe that $E_k \subset \{U_{n_k^*} > x_1, V_{n_k^*} > x_2\}$. Hence the lemma will be established if we show that

$$P(E_k \text{ i.o.}) = 1. \tag{3.1}$$

This in turn will follow when we show as $n \rightarrow \infty$ that

$$E(J_n) \rightarrow \infty \text{ and} \tag{3.2}$$

$$\frac{J_n}{E(J_n)} \rightarrow 1 \text{ in probability} \tag{3.3}$$

where $J_n = \sum_{k=N}^n I_k$ for sufficiently large N , I_k being the indicator function of E_k .

In order to establish (3.2) consider

$$P(E_k) - P(E_k^*) = P(F_k)P(G_k) - P(F_k^*)P(G_k^*) \tag{3.4}$$

$$= P(F_k)\{P(G_k) - P(G_k^*)\} + P(G_k^*)\{P(F_k) - P(F_k^*)\}.$$

Therefore

$$|P(E_k) - P(E_k^*)| \leq P(F_k)|P(G_k) - P(G_k^*)| + P(G_k^*)|P(F_k) - P(F_k^*)| = A_k \text{ say.} \tag{3.5}$$

Observe

From the tail behavior of $\Phi(x)$ (d.f. of standard normal r.v.) that

$$(a) P(E_k) \leq m_k P(X_1 > d_{n_k^*}(x_1)) \sim (\text{const.}) (\log k)^{-1/2} (\log n_k^*)^{-(x_1+1/2)} \text{ as } k \rightarrow \infty.$$

$$(b) P(F_k^*) = 1 - \Phi^{m_k}(d_{n_k^*}(x_1)) = 1 - \exp\left\{m_k \log\left[1 - \left(1 - \Phi(d_{n_k^*}(x_1))\right)\right]\right\}$$

$$= 1 - \exp\left\{-m_k \left(1 - \Phi(d_{n_k^*}(x_1))\right)\right\} (1 + o(1)) \sim (\text{const.}) (\log k)^{-1/2} (\log n_k^*)^{-(x_1+1/2)} \text{ as } k \rightarrow \infty, \text{ whenever } x_1 > -\frac{1}{2}.$$

Similarly $P(G_k^*) \sim (\text{const.}) (\log k)^{-1/2} (\log n_k^*)^{-(x_2+1/2)}$ as $k \rightarrow \infty$, whenever $x_2 > -\frac{1}{2}$.

(2) By lemma 3.1 of Berman (1964)

$$|P(F_k) - P(F_k^*)| = |P(F_k^c) - P(F_k^{*c})| \leq (2\pi)^{-1} \sum_{j=1}^{m_k-1} |r(j)| (m_k - j) (1 - r^2(j))^{-1/2} \times \exp\left\{-d_k^{*2}(x_1)/(1 + |r(j)|)\right\}$$

$$\text{and similarly } |P(G_k) - P(G_k^*)| \leq (2\pi)^{-1} \sum_{j=1}^{m_k-1} |r(j)| (m_k - j) (1 - r^2(j))^{-1/2} \times \exp\left\{-d_k^{*2}(x_2)/(1 + |r(j)|)\right\}$$

By lemma 2.3 and lemma 2.4 we get $\sum_{k=N}^{\infty} A_k < \infty$ whenever either $(\log n)^{1+\gamma} r(n) = O(1)$ as $n \rightarrow \infty$ for some $\gamma > 0$ or $\sum_{j=1}^{\infty} r^2(j) < \infty$.

Further $P(F_k^*)P(G_k^*) \sim (\text{const.}) (\log k)^{-1} (\log n_k^*)^{-(x_1+x_2+1)}$ as $k \rightarrow \infty$.

Since $\limsup \frac{n_k}{n_{k+1}} < 1$, we have $\log n_k^* > (\text{const.}) k^{\frac{1}{1+x_1+x_2}}$ for large k . Hence

$$P(F_k^*)P(G_k^*) \sim (\text{const.}) (\log k)^{-1} k^{-1} \text{ as } \rightarrow \infty. \text{ Thus } \sum P(F_k^*)P(G_k^*) < \infty. \text{ Hence from (3.5), (3.2) follows.}$$

By Chebycheff's inequality we have

$$P\left(\left|\frac{J_n}{E(J_n)} - 1\right| > \varepsilon\right) \leq \frac{V(J_n)}{(\varepsilon^2(EJ_n)^2)} = \frac{\sum_{k=N}^{\infty} V(I_k) + 2 \sum_{N \leq k < l \leq n} \text{Cov}(I_k, I_l)}{\varepsilon^2(EJ_n)^2} \tag{3.6}$$

Clearly $\sum_{k=N}^{\infty} V(I_k) \leq \sum_{k=N}^{\infty} E(I_k) = o(EJ_n)^2$ as $n \rightarrow \infty$. Hence (3.3) will be established if we show

$\sup_n \left| \sum_{N \leq k < l \leq n} \text{Cov}(I_k, I_l) \right| < \infty$ where N is a large positive integer. But this follows from lemma 2.5 and lemma 2.6.

Hence the proof of the lemma is complete.

Lemma 3.3: For all $x_1, x_2 > -\frac{1}{2}$ with $x_1 + x_2 \geq \varepsilon^* - \frac{1}{2}$ and for all $\varepsilon > 0$ we have

$$P(U_{n_k} > x_1 + \varepsilon, V_{n_k} > x_2 + \varepsilon \text{ i.o.}) = 0.$$

Proof: Since U_{n_k} and V_{n_k} are independent, we have

$$P(U_{n_k} > x_1 + \varepsilon, V_{n_k} > x_2 + \varepsilon) = P(U_{n_k} > x_1 + \varepsilon) P(V_{n_k} > x_2 + \varepsilon)$$

$$< n_k^{-2} P(X_1 > d_{n_k}(x_1 + \varepsilon)) P(Y_1 > d_{n_k}(x_2 + \varepsilon))$$

$< (\text{const.}) n_k^2 (d_{n_k}(x_1 + \varepsilon))^{-1} (d_{n_k}(x_2 + \varepsilon))^{-1} \exp\left\{-\frac{1}{2} d_{n_k}^2(x_1 + \varepsilon) - \frac{1}{2} d_{n_k}^2(x_2 + \varepsilon)\right\}$, $k \geq k_{16}$ by the tail behavior of the standard normal distribution.

$< (\text{const.})(\log n_k)^{(1+x_1+x_2+2\varepsilon)}$. Hence $\sum P(U_{n_k} > x_1 + \varepsilon, V_{n_k} > x_2 + \varepsilon) < \infty$

Since $\frac{1}{2} + x_1 + x_2 + 2\varepsilon > \varepsilon^*$ as $x_1 + x_2 \geq \varepsilon^* - \frac{1}{2}$. An application of Borel-Cantelli lemma completes the proof.

Proof of theorem 3.1: From lemmas 2.1 and 2.2 it is clear that the limit set of (U_{n_k}, V_{n_k}) is contained in the square $\{(x_1, x_2) : -\frac{1}{2} \leq x_1, x_2 \leq \varepsilon^*\}$. It follows from lemma 3.3 that the limit set is contained in S_2 .

We conclude from lemmas 3.1 and 3.2 that every point of S_2 except the point $(-\frac{1}{2}, -\frac{1}{2})$ is a limit point. That the point $(-\frac{1}{2}, -\frac{1}{2})$ is also a limit point follows from continuity considerations.

ACKNOWLEDGEMENT

The first author is thankful to the Department of Science and Technology, New Delhi for the financial assistance.

REFERENCES

1. Berman, S.M. (1964): Limit Theorems for the maximum term in stationary sequences. *Ann. Math. Statist.* 35 502-516.
2. Lepage, R.D. (1972/73): Loglog law for Gaussian processes. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* 24 103-108.
3. Mittal, Y. (1974): Limiting behavior of maxima in stationary Gaussian sequences. *Ann. Probability* 2 231-242.
4. Nayak, S.S. (1984): Almost sure limit points of and the number of boundary crossings related to SLLN and LIL for record times, inter-record times and the number of record values. *Stoch. Process. Appl.* 17, 167-176.
5. Nayak, S.S. (1985): Almost sure limit points of independent copies of sample maxima. *Stoch. Process. Appl.* 20, 353-360.
6. Nayak, S.S. (1986): Almost sure limit points maxima of samples from s independent populations. *Sankhya* vol 48, Ser. A Pt.3 273-287.
7. Nayak, S.S. (1988): Almost sure limit points maxima of sample extremes of an IID sequence. *Acta Math. Hung.* 51 (1-2) 35-45.
8. Nayak, S.S. and Madhusudhan Zalki (2000): On the fluctuations of independent copies of moving maxima. *Statistics and Probability Letters*. vol.50 351-356.
9. Nayak, S.S. and Madhusudhan Zalki (2001): Almost sure limit points of record values from two independent populations. *Statistics and Probability Letters*. vol.51 181-187.
10. Nayak, S.S. and WALL, K.S. (1994): Almost sure limit points maxima of sample extremes of Samples from two independent populations. *Sankhya* vol. 56 Ser.A Pt.2 294-319.
11. Pakshirajan, R.P. and Vasudeva, R. (1977): A law of the iterated logarithm for stable summands. *Trans. Amer. Math. Soc.* 232 33-42.
12. Pickands, J. III (1969): An iterated logarithm law for the maximum in a stationary Gaussian sequence. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* 12 344-353.
13. Qualls, C. and Watanbe, H. (1971): An asymptotic 0-1 behaviour of Gaussian processes. *Ann. Math. Statist.* 42 2029-2035.
14. Strassen, V. (1964): An invariance principle for the law of the iterated logarithm. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* 3 211-226.
15. Vasudeva R. and Savita S. (1995): Law of the iterated logarithm for subsequences of partial maxima of stationary Gaussian sequences. *Journal of the Indian Society for Probability and statistics* 2 15-30.
16. Vishnu Hebbar, H. (1980): Almost sure limit points of maxima of stationary Gaussian sequences. *Ann. Probability* 8 No.2 393-399.

Source of Support: None Declared
Conflict of Interest: None Declared